



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

DNA methylation profiling to predict recurrence risk in meningioma: development and validation of a nomogram to optimize clinical management

Nassiri, Farshad ; Mamatjan, Yasin ; Suppiah, Suganth ; Weller, Michael ; et al

Abstract: BACKGROUND Variability in standard-of-care classifications precludes accurate predictions of early tumor recurrence for individual patients with meningioma, limiting the appropriate selection of patients who would benefit from adjuvant radiotherapy to delay recurrence. We aimed to develop an individualized prediction model of early recurrence risk combining clinical and molecular factors in meningioma. METHODS DNA methylation profiles of clinically annotated tumor samples across multiple institutions were used to develop a methylome model of 5-year recurrence-free survival (RFS). Subsequently, a 5-year meningioma recurrence score was generated using a nomogram that integrated the methylome model with established prognostic clinical factors. Performance of both models was evaluated and compared with standard-of-care models using multiple independent cohorts. RESULTS The methylome-based predictor of 5-year RFS performed favorably compared with a grade-based predictor when tested using the 3 validation cohorts ($\Delta\text{AUC} = 0.10$, 95% CI: 0.03-0.018) and was independently associated with RFS after adjusting for histopathologic grade, extent of resection, and burden of copy number alterations (hazard ratio 3.6, 95% CI: 1.8-7.2, $P < 0.001$). A nomogram combining the methylome predictor with clinical factors demonstrated greater discrimination than a nomogram using clinical factors alone in 2 independent validation cohorts ($\Delta\text{AUC} = 0.25$, 95% CI: 0.22-0.27) and resulted in 2 groups with distinct recurrence patterns (hazard ratio 7.7, 95% CI: 5.3-11.1, $P < 0.001$) with clinical implications. CONCLUSIONS The models developed and validated in this study provide important prognostic information not captured by previously established clinical and molecular factors which could be used to individualize decisions regarding postoperative therapeutic interventions, in particular whether to treat patients with adjuvant radiotherapy versus observation alone.

DOI: <https://doi.org/10.1093/neuonc/noz061>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-176277>

Journal Article

Accepted Version

Originally published at:

Nassiri, Farshad; Mamatjan, Yasin; Suppiah, Suganth; Weller, Michael; et al (2019). DNA methylation profiling to predict recurrence risk in meningioma: development and validation of a nomogram to optimize clinical management. *Neuro-Oncology*, 21(7):901-910.

DOI: <https://doi.org/10.1093/neuonc/noz061>

DNA methylation profiling to predict recurrence risk in meningioma: development and validation of a nomogram to optimize clinical management

Farshad Nassiri MD^{1,2*}, Yasin Mamatjan PhD^{2*}, Suganth Suppiah MD^{1,2}, Jetan H Badhiwala MD², Sheila Mansouri PhD², Shirin Karimi MD², Olli Saarela PhD³, Laila Poisson PhD⁴, Irina Gepfner-Tuma MD⁵, Jens Schittenhelm MD⁶, Ho-Keung Ng MD⁷, Houtan Noushmehr PhD⁸, Patrick Harter MD⁹, Peter Baumgarten MD¹⁰, Michael Weller MD¹¹, Matthias Preusser MD¹², Christel Herold-Mende PhD¹³, Marcos Tatagiba MD¹⁴, Ghazaleh Tabatabai MD⁵, Felix Sahm MD PhD¹⁵ PhD, Andreas von Deimling MD PhD¹⁵, Gelareh Zadeh MD PhD^{1,2,16,18**}, Kenneth D Aldape MD^{1,2,17,18**}, and the International Consortium on Meningiomas⁺

*contributed equally to this work

**co-corresponding authors

+see authorship statement at end of article

¹ Division of Neurosurgery, Department of Surgery, University of Toronto, Toronto, Ontario, Canada

² MacFeeters-Hamilton Centre for Neuro-Oncology Research, Princess Margaret Cancer Centre, Toronto, Ontario, Canada

³ Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario, Canada

⁴ Department of Public Health Sciences, Henry Ford Health System, Detroit, MI, USA

⁵ Interdisciplinary Division of Neuro-Oncology, Hertie Institute for Clinical Brain Research & Centre for CNS Tumors, Comprehensive Cancer Center Tübingen-Stuttgart, University Hospital Tübingen, Eberhard Karls University, Tübingen, Germany

⁶ Department of Neuropathology, Institute of Pathology and Neuropathology, University of Tübingen, Tübingen, Germany

⁷ Department of Anatomical and Cellular Pathology, The Chinese University of Hong Kong, Hong Kong, China

⁸ Department of Neurosurgery, Henry Ford Health System, Detroit, MI 48202, USA

⁹ Edinger Institute (Institute of Neurology), Goethe-University, Frankfurt am Main, Germany

¹⁰ Department of Neurosurgery, Goethe-University, Frankfurt am Main, Germany

¹¹ Department of Neurology, University Hospital Zurich, Zurich, Switzerland

¹² Clinical Division of Oncology, Department of Medicine I, Comprehensive Cancer Centre CNS Unit, Medical University of Vienna, Vienna, Austria

¹³ Department of Neurosurgery, Heidelberg University Hospital, Heidelberg, Germany

¹⁴ Department of Neurosurgery, University Hospital Tübingen, Tübingen, Germany

¹⁵ Department of Neuropathology, Institute of Pathology, Ruprecht-Karls-University Heidelberg, Heidelberg, Germany

¹⁶ Division of Neurosurgery, University Health Network, Toronto, Ontario, Canada

¹⁷ Laboratory of Pathology, Centre for Cancer Research, National Cancer Institute, Bethesda, MD, USA

¹⁸ These authors jointly directed this work. Correspondences should be addressed to K.D.A, National Cancer Institute, Building 10, Room 2S235, Bethesda, MD 20892-1500, Ph: 301-480-8853 (kenneth.aldape@nih.gov) or G.Z, Princess Margaret Cancer Research Tower, 101 College Street, 14th floor, Room 601, M5G 1L7, (gelareh.zadeh@uhn.ca)

Running head: Individualized predictor of recurrence in meningiomas

Conflict of Interest: The authors have no personal, financial, or institutional interest in any of the drugs, materials, or devices described in this article.

Funding: The Brain Tumor Charity Quest for Cures: Collaborative Team Award, Canadian Institute of Health Research - Institute of Cancer Research Operating Grant

Authorship: KA, GZ, FN conceived, designed, and supervised the study. GZ, KA, FS, PH, MW, MP, CHH, AvD, IGT, HKN, MT, GT collected the specimens. FN, MY, SS, SK, SM, JHB, constructed the database. KA, FS, AvD, HN, JS offered pathological review. MY, FN LP, OS did the statistical and computational analyses. FN, MY, KA, GZ wrote the manuscript and all authors revised the manuscript. All authors discussed and reviewed the manuscript and approved the manuscript for publication.

Word Count: 5618

CONFIDENTIAL

ABSTRACT

Background: Variability in standard-of-care classifications preclude accurate predictions of early tumor recurrence for individual patients with meningioma limiting the appropriate selection of patients who would benefit from adjuvant radiotherapy to delay recurrence. We aimed to develop an individualized prediction model of early recurrence risk combining clinical and molecular factors in meningioma.

Methods: DNA methylation profiles of clinically-annotated tumor samples across multiple institutions were used to develop a methylome-model of 5-year recurrence-free-survival (RFS). Subsequently, a 5-year meningioma recurrence score was generated using a nomogram that integrated the methylome-model with established prognostic clinical factors. Performance of both models was evaluated and compared to standard-of-care models using multiple independent cohorts.

Results: The methylome-based predictor of 5-year RFS performed favorably compared to a grade-based predictor when tested using the three validation cohorts ($\Delta\text{AUC} = 0.10$, 95%CI 0.03–0.018) and was independently associated with RFS after adjusting for histopathological grade, extent of resection and burden of copy number alterations (HR 3.6, 95%CI 1.8–7.2, $P < 0.001$). A nomogram combining the methylome-predictor with clinical factors demonstrated greater discrimination than a nomogram using clinical factors alone in two independent validation cohorts ($\Delta\text{AUC} = 0.25$, 95%CI 0.22 – 0.27) and resulted in two groups with distinct recurrence patterns (HR 7.7, 95%CI 5.3–11.1, $P < 0.001$) with clinical implications.

Conclusions: The models developed and validated in this study provide important prognostic information not captured by previously established clinical and molecular

factors which could be used to individualize decisions regarding post-operative therapeutic interventions, in particular, whether to treat patients with adjuvant radiotherapy versus observation alone

Keywords: methylation, meningioma, nomogram, predictor, recurrence

Key points:

- Combining DNA methylation with clinical factors results in reliable individualized estimations of recurrence risk
- Individualized recurrence risk can be used to guide post-operative therapeutic interventions

Importance of the Study

Our work is the first to demonstrate the transformative utility of integrating clinical and molecular factors for use beyond simple classification into the realm of individualized prognostication for any brain tumor. Using our developed and validated tools that are publicly available, clinicians will be able combine clinical and molecular factors to determine an **individualized** probability of recurrence for patients with meningiomas. This represents a major advance in the field of personalized medicine for neuro-oncology, and the use of this tool can help clinicians overcome one of the most challenging limitations we face when treating patients with meningiomas.

INTRODUCTION

Meningiomas are the most common primary intracranial tumor. They account for 37% of all central nervous system tumors, and are continuing to increase in incidence with the ageing population¹. They result in significant neurological morbidity and loss of quality of life by exerting mass effect on critical adjacent brain regions¹. The current standard-of-care for nearly all patients with symptomatic meningiomas includes gross total tumor resection with removal of involved dura and bone when possible². However, despite radical surgical resection, approximately 20% of meningiomas display aggressive behaviour with early tumor recurrence resulting in a clinical course of repetitive disease- and treatment-related morbidity. Radiation therapy can be used to provide disease control as an adjunct to surgery for a subset of tumors^{3,4}. However, radiotherapy can often result in adverse radiation effects that lead to considerable morbidity and neurological dysfunction long-term precluding universal use in all patients⁵. One of the greatest clinical challenges faced by clinicians is the inability to predict early tumor recurrence at an individual patient level which limits the appropriate selection of patients who would benefit from adjuvant radiation therapy.

To date, the most reliable clinical factors associated with recurrence in meningiomas have been the World Health Organization (WHO) grade of the tumor and extent of tumor resection at surgery^{2,6}. Although both are crudely associated with recurrence rates on a population level, they are challenged with inter-rater variability of grading and considerable within-grade variation of recurrence risk for individual patients^{2,6}. In the past decade, several studies have focused on molecular profiling of meningiomas to refine biological subgroups⁷⁻¹³. With the exception of mutations in *BAP1* and *TERT* promoter,

each of which occur rarely in these tumors, the mutations identified in meningioma have not been shown to be tightly correlated to patient outcome with current standard of care⁷⁻¹⁰. We and others have independently shown that global DNA methylation profiling reveals robust methylome-based meningioma subtypes, however, the clinical translation of this to predict recurrence risk for individual patients has not been demonstrated to date.

To examine whether methylation profiles can be defined and validated for clinical utility, we aimed to develop and validate a methylome-based predictor of early meningioma recurrence that could be combined with established prognostic clinical factors to individualize decisions regarding the need for post-operative therapeutic interventions, in particular, whether to treat patients with adjuvant radiation therapy versus observation alone. Our work is the first in neuro-oncology to demonstrate the transformative utility of integrating clinical and molecular factors for use beyond simple classification into the realm of individualized prognostication.

MATERIALS AND METHODS

Study design and data sources

This multicenter retrospective study was carried out in accordance with individual institutional ethics and review board guidelines and comprised a total of 486 patients with clinically annotated and available meningioma samples. Institutional waivers of informed consent were obtained due to minimal patient risk associated with this study. Two-hundred and eighty-two (N=282) fresh-frozen or formalin-fixed paraffin embedded (FFPE) meningioma tumor samples from multiple institutions (N=76 from Princess Margaret Cancer Research Centre, Toronto, Canada; N=206 from European centers including

University of Heidelberg, Heidelberg, Germany and Goethe-University, Frankfurt, Germany and University of Tübingen, Tübingen, Germany, and University Hospital Zurich, Zurich, Switzerland and Medical University of Vienna, Vienna, Austria) comprised the discovery cohort. which was split into a training cohort (81%, N=228 samples) and first validation cohort (19%, N=54 samples) each balanced for tumor grade, tissue type, recurrence status and time to recurrence. One hundred and forty (N=140) FFPE meningioma tumor samples from a separate institution (MD Anderson Cancer Centre, Houston, USA) were used as a second validation cohort, and sixty-four (N=64) fresh frozen meningioma tumor samples from two other institutions were used as a third validation cohort (N=46 from Princess Margaret Cancer Research Centre; N=18 from The Chinese University of Hong Kong). The sample sets from Europe and MD Anderson comprised the subset of previously published samples for which clinical data (recurrence-free survival [RFS], WHO grade) were available^{11,12}. Moreover, *TERT* promoter mutation status was available on a subset of previously published European samples^{10,11}. Gene expression analysis was performed on publicly available microarray data on 98 patients with meningiomas of all grades (GSE16581¹⁴ and GSE9438¹⁵). The outline for the overall study design is demonstrated in Supplementary Figure S1.

Definitions

Hematoxylin and eosin (H&E) slides for each patient were reviewed for meningioma diagnosis and WHO grading was performed according to the current WHO 2016 criteria at local institutions by experienced neuropathologists. Tumor recurrence and time to recurrence were the primary outcomes of interest and were collected locally for each

sample as previously described^{11,12}. Briefly, recurrence was defined as tumor growth following gross-total resection or tumor progression following subtotal resection. Time to recurrence was determined by reviewing post-operative imaging and calculating the duration from the date of surgery to first post-operative imaging documenting tumor recurrence in concordance with documentation in the medical charts. The extent of resection (Simpson grade²) was determined based on the surgeon's operative report in correlation with postoperative cranial imaging.

Generation of an individualized methylome-predictor of 5-year RFS

For DNA methylation and copy number analysis of the samples, DNA was extracted from each tumor and DNA methylation profiling was performed using Illumina 450k HumanMethylation BeadChip or 850k EPIC arrays as per manufacturer instructions at each institution. Raw data files (*.idat) were imported, processed, and normalized to integrate data from multiple generations of Infinium methylation arrays. Copy number aberrations were inferred from methylation array data¹⁶ and burden of copy number alterations was computed per sample as previously described¹⁷. Probes that were common in both 450k HumanMethylation Beadchip and 850K EPIC arrays were selected as possible features for development of our predictor such that our predictor would be applicable to the landscape of available technologies. We used a multi-step strategy to select the probes to be used in the generation of our predictor (See Supplementary Methods and Supplementary Figure S2).

To develop the methylome-based predictor of early meningioma recurrence, we performed generalized boosted regression modelling using the final selected probes in samples from the training cohort to predict 5-year RFS. Boosted regression modelling using WHO grade as a sole feature in the training cohort was also performed and tested in each validation cohort to compare methylation-based predictor performance to a standard-of-care model. Performance of both models was assessed by generating time-dependent receiver operating characteristic (ROC) curves and computing average areas under the ROC curves (AUC) for each validation cohort independently, along with their 95% confidence intervals using the bootstrap resampling method with 10,000 resamples¹⁸.

Methylation probe annotation was performed using the UCSC Genome Browser (GRCh38/hg38 assembly). We used the Functional Annotation Clustering algorithm¹⁹ of Database for Annotation, Visualization, and Integrated Discovery (DAVID)¹⁹ Bioinformatics Resource 6.8 to identify redundant functional clusters represented by genes annotated with a minimum of 5 probes (See Supplementary Figure S3). Two publicly available microarray datasets (GSE16581¹⁴ and GSE9438¹⁵) reporting on 22486 genes for 98 patients with meningiomas were pooled to correlate methylation data with gene expression data as an exploratory analysis.

Further details regarding the steps for generation, validation and characterization of the methylome-predictor of 5-year RFS can be found in Supplementary Methods.

Generation of a meningioma-recurrence score

To create a contemporary meningioma recurrence score that could be utilized by clinicians to predict early risk of recurrence for individual patients, we generated a nomogram based on a Cox model that incorporated the methylome-based predictor, WHO grade, and extent of resection using samples from the training cohort and second validation cohort to predict 5-year RFS. Both the training cohort and second validation cohort were used to train this nomogram in order to increase the number of samples available to capture the heterogeneity in the spectrum of data available. It is important to note that as none of the samples used to train this model were used to assess model performance on external validation. The global performance of the meningioma recurrence was assessed by generating time-dependent receiver operating characteristic (ROC) curves and computing average area under the ROC curves (AUC) for the first and third validation cohort independently, along with their 95% confidence intervals using the bootstrap resampling method with 10,000 resamples^{18,20}. For comparison, a nomogram that incorporated only WHO grade and extent of resection as the sole features was also developed in similar fashion. Internal validation using bootstrap resampling using 10,000 resamples was also performed. Model calibration was assessed visually by plotting observed event rates against nomogram predicted probabilities for two risk groups. Detailed descriptions on nomogram calculations and use are described in the Supplementary Methods and Supplementary Figure S4.

Statistical analysis

Summary statistics are reported as counts (and proportions) for categorical variables and median (and range) for continuous variables, unless otherwise indicated. Cohort size was

determined by availability of samples. Statistical analyses were performed in consultation with two expert biostatisticians (L.P and O.S).

To investigate the clinical relevance of the methylome-based predictor and meningioma recurrence score, distribution of survival times was performed using Kaplan-Meier methods and compared across groups using log-rank testing. The frequency of genome-wide copy number alterations across groups was computed and plotted using a custom algorithm. The performance of a methylome-based predictor was compared to grade-based predictor by computing an average ΔAUC ($AUC_{\text{methylome}} - AUC_{\text{grade}}$) and 95%CI from all bootstraps and the performance of the meningioma recurrence score incorporating the methylome predictor was compared to a nomogram excluding the methylome predictor by computing average ΔAUC ($AUC_{\text{combined nomogram}} - AUC_{\text{clinical only nomogram}}$) and 95%CI in similar fashion.

Hazard ratios, including 95% confidence intervals (CI), were calculated based on univariable and multivariable Cox regression modeling for the methylome-based predictor and other covariates including WHO grade, extent of resection, and burden of copy number alterations. Proportional hazards assumption was tested by computing Schoenfeld residuals for each covariate and testing was performed according to Grambsch and Therneau²¹ (Supplementary Figure S5 and Supplementary Table S1). Martingale residuals were plotted against methylation predictor probabilities to determine appropriateness of linear modeling (Supplementary Figure S6). Sensitivity analyses evaluating the possible confounding effects of TERT promoter mutations, receipt of adjuvant radiation therapy,

and center-effects were performed using multivariable Cox regression in samples with available information. Comparison of proportions across groups was completed using χ^2 test and Fisher Exact test, where appropriate.

Two-sided p-values are reported, and the threshold for statistical significance was set a priori at $\alpha = 0.05$. We used R version 3.3.1 for all statistical analyses, model generation, and model validation.

RESULTS

A set of 9529 probes were selected from an initial training cohort and generalized boosted regression modeling was performed to develop a DNA methylation-based predictor of 5-year recurrence risk in meningioma (Supplementary Figure S1). Three validation cohorts (Supplementary Table S2) were used to test the performance of the methylation-based predictor compared to a grade-based predictor.

Validation of a methylome-predictor of early meningioma recurrence

The methylome-based predictor performed favorably compared to the grade-based predictor at 5-years in each of the 3 validation cohorts (Figure 1) with substantial statistical improvements in performance when tested in all combined validation cohorts ($\Delta\text{AUC} = 0.10$, 95%CI 0.03 – 0.18). When stratified by median, the 5-year methylome-predictor distinguished risk groups (lower and higher risk) in all 3 validation cohorts (Figure 2). Patients in the higher risk group had a median RFS of 2.1 years, 8.1 years, and 4.2 years in the first, second, and third validation cohorts respectively compared to patients

in the lower risk groups which had median RFS of “unreached” in the first and second validation cohorts and median RFS of 7.2 years in the third validation cohort (HR 8.1 95%CI 2.8 – 23.8, HR 3.5 95%CI 1.8 – 6.6, and HR 2.0, 95%CI 1.2 – 3.7, respectively).

DNA copy number analysis demonstrated that increased frequency of copy number aberrations in the higher risk groups, as demonstrated by high proportion of chromosomal deletions in 1p, 4p, 6q, 10q, 14q, and 18q (Figure 3). The total burden of copy number alterations was also correlated with risk groups, with greatest proportion of burden of copy number alterations found in the higher risk groups in all three validation cohorts (Supplementary 03). It is noteworthy that of all patients with high burden of copy number aberrations from all three validation cohorts, only 12 patients (4.6%) were in the lower risk group.

Multivariable Cox regression analysis demonstrated that the 5-year methylome based predictor was independently associated with RFS in samples from all validation cohorts (HR 3.6, 95%CI 1.8 – 7.2, $P < 0.001$) after controlling for tumor grade, extent of resection, burden of copy number alterations (Table S4). Sensitivity analyses including receipt of adjuvant therapy, TERT promoter mutations, and center of treatment as covariates did not alter this relationship and these covariates were not independently associated with RFS (Supplementary Tables S5-7).

Current clinical practice relies on histopathologic grade to inform the decision of post-surgical management of meningioma. Patients with grade I tumors are most commonly

monitored with serial imaging after surgery, while patients with grade II and III tumors are considered for adjuvant therapy, such as radiation, to prevent recurrence. There is, however, unexplained clinical variability in RFS within all grades of meningioma, and we examined the use of the 5-year methylome-predictor to address this issue. Among patients with WHO grade II tumors in all validation cohorts, the median RFS for the higher risk group was 2.6 years compared to a median RFS of 8.4 years in the lower risk group (HR 2.8, 95%CI 1.7 – 4.8, $P < 0.001$, Figure 4B). Patients with WHO Grade I tumors in all validation cohorts also had increased risk for recurrence in the higher risk group (HR 2.9, 95%CI 1.3 – 6.6, $P=0.006$, Figure 4A) with only 3 patients in the lower risk group recurring within the first 5 years compared to 18 patients in the higher risk group. Lastly, patients with WHO Grade III tumors from all validation cohorts had poor median RFS in the higher risk group (1.3 years), compared to a median RFS of 6.0 years in the lower risk group (HR 3.0, 95%CI 1.4 – 6.5, $P=0.004$, Figure 4C).

Characterization of predictor CpG sites

The selected 9529 probes used in our model comprise only 1% of all probes included on the 850K Illumina Array. These probes were enriched to be found in the promoter regions ($N=3057$, 32.1%) and located on CpG islands ($N = 4633$, 48.6%) compared to all probes found on the 850K Illumina Array (29.6%, $P < 0.001$ and 18.0%, $P < 0.001$, respectively; Supplementary Table S8) Of the 9529 probes, only 1261 (13.2%) were “favorable” probes associated with lower risk of recurrence when methylated (HR ranging from 0.002 to 0.38, on univariable Cox regression with associated $P < 0.001$). The remaining 8237 (86.8%) probes were “unfavorable” probes associated with higher risk of recurrence when

methyated (HR ranging from 2.45 to 517.82, on univariable Cox regression with associated $P < 0.001$).

There were 2332 probes annotated to 294 genes with at least 5 probes represented per gene. Of these (Supplementary Table S9), only 68 of 2332 probes (2.9%) were found to be “favorable” probes (associated with lower recurrence risk when methyated, HR ranging from 0.008 to 0.252 on univariable Cox regression with associated $P < 0.001$). The remaining 2265 probes (97.1%) were found to be “unfavorable” probes associated with greater recurrence risk (associated with higher recurrence risk when methyated, HR ranging from 2.95 to 517.82 on univariable Cox regression with associated $P < 0.001$). Functional annotation clustering of these 2265 “unfavorable” probes revealed that homeobox (enrichment score = 54.33) and TBox (enrichment score = 4.12) were highly significant redundant functional clusters (Supplementary Figure S3). Gene expression analysis of Homeobox family of genes and Tbox genes for which methylation data was also available, demonstrated that although these genes were relatively hypermethyated in recurrence-prone tumors, they were either upregulated or non-differentially expressed (Supplementary Table S10).

Validation and clinical utility of a meningioma recurrence score

To generate and validate a meningioma recurrence score that could be translated to the clinic, we developed a nomogram to predict 5-year RFS that incorporated the validated 5y methylome-based predictor with established prognostic covariates (WHO grade, and extent of resection) using samples from the training cohort and second validation cohort

(Figure 5). Validation of the nomogram using the first validation cohort and third validation cohort independently as well as both validation cohorts combined demonstrated greater predictive performance of the nomogram with methylome predictor included compared to a nomogram using clinical factors (WHO grade and extent of resection) alone ($\Delta\text{AUC} = 0.24$, 95%CI 0.19 – 0.28; $\Delta\text{AUC} = 0.26$, 95%CI 0.22 – 0.31; $\Delta\text{AUC} = 0.25$, 95%CI 0.22 – 0.27 respectively, Figure 5). The discriminative power of the meningioma recurrence score was approximately 82% in combined validation cohorts ($\text{AUC} = 0.82$, 95%CI 0.76 – 0.87). The performance of the model using external validation and internal validation approaches were similar (Supplementary Figure S7). The highest proportion of allottable points given in the nomogram is based on probabilities from the methylome-based predictor, again suggesting that the methylome-predictor has greater importance in determining recurrence risk in meningiomas compared to established clinical factors. Calibration and Kaplan-Meier survival analysis of the meningioma recurrence score clearly stratifies patients with high risk and low risk for 5-year recurrence (HR 7.7, 95%CI 5.3 – 11.1, $P < 0.001$, Figure 6). Interestingly, while histologic grades II and III are meant to predict high recurrence risk, we find that the low-risk group in total contains 39 grade II tumors (34.5% of the grade II tumors) and 4 grade III tumors (8.3% of the grade III tumors). Conversely, while a WHO grade I designation is meant to convey a low risk of recurrence, there were 35 (21.2%) patients with grade I tumors in the high-risk group. These results indicate refinement of risk estimate by the nomogram relative to current classification standards. To facilitate unrestricted global dissemination, we have created a freely available online calculator of the meningioma recurrence score https://meningiomas.shinyapps.io/meningioma_recurrence_score_online_calculator/.

Details on imputations for the online calculator can be found in the Supplementary Methods, and case examples demonstrating the power of personalized predictions are detailed in Supplementary Figures S8-10.

DISCUSSION

In this multicenter study, we demonstrated the transformative utility of integrating clinical and molecular factors for use beyond simple classification into the realm of individualized prognostication in neuro-oncology. Our methylome-based predictor was more reliably able to predict early (5-year) recurrence-free survival in comparison with histologic grading and was associated with RFS independent of established clinical and molecular factors. Combining the methylome predictor with established prognostic clinical factors (WHO grade and extent of resection) in a meningioma recurrence score refined prognostication for individual patients with meningiomas beyond established prognostic clinical and molecular factors with therapeutic implications for individualizing decision-making regarding the need for adjuvant therapy after surgery in meningiomas.

Although WHO grade is associated with recurrence in populations of meningioma patients, and is currently used to guide therapy, the clear within-grade variation for risk of recurrence and inter-rater variability makes it challenging to rely on tumor grade alone to predict recurrence and guide postoperative management decisions for individual patients. As a manifestation of this imprecision, some patients with biologically aggressive tumors may be inappropriately subsumed in within the group of histologically benign tumors. With current standard of care, it is thought that up-front treatment with adjuvant radiation therapy after surgical resection offers the best chance to delay recurrence, and therefore

some patients are not appropriately selected for adjuvant treatment with standard-of-care approaches^{3,4}. Conversely, there are also some patients with histologically defined intermediate or higher-grade tumors that in fact harbor indolent tumor behavior. Such patients may be receiving adjuvant radiation therapy in the absence of a defined need. Radiation, even when optimized to minimize adverse effects, still carries the risk for adverse radiation effects such as reactive inflammation, vasculitis, and necrosis, all of which have sequelae on patient cognition and quality of life⁵. There is a clear need for a more refined predictor of recurrence patterns for individual patients with meningiomas beyond simple classifications, so that the decision for adjuvant therapy, radiotherapy or otherwise, can be appropriately selected and personalized for patients.

The burden of chromosomal alterations have repeatedly been shown to be one of the most important prognostic molecular alterations in meningioma^{11,17,22}. Similar to others^{11,22}, we found recurrent alterations in 1p, 4p, 6q, 10q, 14q, and 18q in a subset of meningiomas that were enriched in higher risk group. However, using a previously published copy number score designed to identify meningiomas with high recurrence risk¹⁷, we found that the burden of copy number alteration was not an independent predictor of recurrence once adjusted for methylation signature. Similarly, although TERT promoter mutations are known to be enriched in more aggressive meningiomas¹⁰, our analysis demonstrated that TERT promoter mutation was not independently associated with RFS on multivariable analysis with the 5-year methylome predictor included in the model. Taken together, these results suggest that our 5-year methylome predictor can provide prognostic information beyond previously established molecular factors in meningioma.

Probes included in the predictor were selected based on correlations of either hypo- or hypermethylated status with respect to RFS. Interestingly, the distribution of these probes was not random, either with respect to methylation status nor with respect to association with known genes. The majority (over 86%) of our included probes were all associated with unfavorable RFS when hypermethylated. This suggests that in general, hypermethylation of a set of CpG sites in meningioma correlates with clinical aggressiveness. For example, hypermethylation of the homeobox and Tbox families of genes were found to be highly overrepresented in the set of relevant probes, suggesting possible involvement of this class of developmental factors in the clinical behavior of meningioma, which requires further validation.

To generate a tool that could be used by clinicians to capture the heterogeneity in recurrence risk in meningiomas, we established a 5-year meningioma recurrence score that combined our validated methylome-based predictor with well-established prognostic clinical factors (WHO grade and extent of resection). The performance of our meningioma recurrence score was improved with the methylome-based predictor included in the nomogram ($\Delta\text{AUC} = 0.25$, 95%CI 0.22 – 0.27), and the overall discrimination of our nomogram was high (AUC approximately 82% in two independent cohorts). The construction, and evaluation, of our nomogram meets the standards of reporting on nomograms in oncology and is one of few to demonstrate robust evaluation using multiple independent validation cohorts²³. Now that we have demonstrated that our tools are robustly validated, we are well positioned to prospectively validate the use of our tools to

demonstrate efficacy with adjuvant therapy strategies in high-risk patients²⁴. Our meningioma recurrence score informs both patients and clinicians about individualized risk of recurrence, can be used to guide clinicians regarding the need for adjuvant therapy and/or close clinical follow-up.

Our study has some limitations. First, although we have identified a group of probes with distinct epigenetic changes that in combination are predictive of recurrence risk in meningiomas for individual patients, it is unclear whether these changes may be conferring variant behavioural phenotypes or whether they are a surrogate for general cellular dysregulation. Nevertheless, the set of highly refined and selected probes in our predictor are enriched to be located on promoters of CpG islands where aberrant DNA methylation has clearly been linked to carcinogenesis²⁵. The correlation of methylation with gene expression in our study was exploratory and would benefit from additional investigation with matched epigenetic and transcriptomic analysis in the same samples. Moreover, although each institution conformed to a common definition for tumour recurrence and time to recurrence, there is no universally standardized definition of recurrence in meningiomas. The Response Assessment in Neuro-Oncology guidelines may offer an avenue for standardized definition of recurrence to be collected on meningiomas in future clinical trials, which will help with communication across different centres and may also help with further model refinement²⁶.

Our predictor has been designed and validated such that it can be applied to data from fresh frozen or paraffin embedded tissues using the current commonly-used standard

platform for genome-wide DNA methylation profiling, facilitating immediate adoption into clinical practice. Our newly developed and retrospectively validated meningioma recurrence score combines both methylome and clinical factors and can be freely used by clinicians to personalize decision making regarding post-operative management of the most common primary intracranial brain tumor via a web-based interface.

DATA ACCESSION: Previously unpublished raw methylation files (idat files) have been submitted to the European Genome-phenome Archive (EGAS00001003490).

AUTHORSHIP STATEMENT: The generation of this manuscript has been supported by the membership of the consortium, which at the time of supplement generation includes: Kenneth Aldape, Karolyn Au, Jill Barnhartz-Sloan, Wenya Linda Bi, Priscilla K. Brastianos, Nicholas Butowski, Carlos Carlotti, Michael D. Cusimano, Francesco DiMeco, Katharine Drummond, Ian F. Dunn, Evanthia Galanis, Caterina Giannini, Roland Goldbrunner, Brent Griffith, Rintaro Hashizume, C. Oliver Hanemann, Christel Herold-Mende, Craig Horbinski, Raymond Y. Huang, David James, Michael D. Jenkinson, Christine Jungk, Timothy J. Kaufman, Boris Krischek, Daniel Lachance, Christian Lafougère, Ian Lee, Jeff C. Liu, Yasin Mamatjan, Tathiane M. Malta, Christian Mawrin, Michael McDermott, David Munoz, Farshad Nassiri, Houtan Noushmehr, Ho-Keung Ng, Arie Perry, Farhad Pirouzmmand, Laila M Poisson, Bianca Pollo, David Raleigh, Felix Sahm, Andrea Saladino, Thomas Santarius, Christian Schichor, David Schultz, Nils O. Schmidt, Warren Selman, Andrew Sloan, Julian Spears, James Snyder, Suganth Suppiah, Ghazaleh Tabatabai, Marcos Tatagiba, Daniela Tirapelli,

Joerg C. Tonn, Derek Tsang, Michael A. Vogelbaum, Andreas von Deimling, Patrick Y. Wen, Tobias Walbert, Manfred Westphal, Adriana M. Workewych, Gelareh Zadeh.

FUNDING: The Brain Tumor Charity Quest for Cures: Collaborative Team Award, Canadian Institute of Health Research - Institute of Cancer Research Operating Grant

ACKNOWLEDGEMENTS: Adriana Workewych for her help in formatting the figures of this manuscript.

REFERENCES

1. Ostrom QT, Gittleman H, Xu J, et al. CBTRUS statistical report: Primary brain and other central nervous system tumors diagnosed in the United States in 2009-2013. *Neuro Oncol.* 2016;18(suppl_5):v1-v75. doi:10.1093/neuonc/nov189.
2. Melamed S, Sahar A, Beller AJ. The recurrence of intracranial meningiomas. *Neurochirurgia (Stuttg).* 1979;22(2):47-51. doi:10.1055/s-0028-1090287.
3. Weber DC, Ares C, Villa S, et al. Adjuvant postoperative high-dose radiotherapy for atypical and malignant meningioma: A phase-II parallel non-randomized and observation study (EORTC 22042-26042). *Radiother Oncol.* 2018;128(2):260-265. doi:10.1016/j.radonc.2018.06.018.
4. Rogers L, Zhang P, Vogelbaum MA, et al. Intermediate-risk meningioma: initial outcomes from NRG Oncology RTOG 0539. *J Neurosurg.* 2018;129(1):35-47.

- doi:10.3171/2016.11.JNS161170.
5. Rogers L, Barani I, Chamberlain M, et al. Meningiomas: knowledge base, treatment outcomes, and uncertainties. A RANO review. *J Neurosurg.* 2015;122(1):4-23.
doi:10.3171/2014.7.JNS131644.
 6. Toro-Moreno AC, Serna-Velez L, Gallego-González D, Jaramillo-Jaramillo LI, Martínez-Sánchez LM, Álvarez-Hernández LF. Tumores de sistema nervioso central en pediatría: Presente y futuro del abordaje diagnóstico. *Rev Ecuatoriana Neurol.* 2017;26(3):283-288. doi:10.1007/s00401-016-1545-1.
 7. Brastianos PK, Horowitz PM, Santagata S, et al. Genomic sequencing of meningiomas identifies oncogenic SMO and AKT1 mutations. *Nat Genet.* 2013;45(3):285-289.
doi:10.1038/ng.2526.
 8. Clark VE, Erson-Omay EZ, Serin A, et al. Genomic analysis of non-NF2 meningiomas reveals mutations in TRAF7, KLF4, AKT1, and SMO. *Science (80-).* 2013;339(6123):1077-1080. doi:10.1126/science.1233009.
 9. Clark VE, Harmancl AS, Bai H, et al. Recurrent somatic mutations in POLR2A define a distinct subset of meningiomas. *Nat Genet.* 2016;48(10):1253-1259.
doi:10.1038/ng.3651.
 10. Sahm F, Schrimpf D, Olar A, et al. TERT Promoter Mutations and Risk of Recurrence in Meningioma. *J Natl Cancer Inst.* 2016;108(5):djv377. doi:10.1093/jnci/djv377.
 11. Sahm F, Schrimpf D, Stichel D, et al. DNA methylation-based classification and grading system for meningioma: a multicentre, retrospective analysis. *Lancet Oncol.* 2017;18(5):682-694. doi:10.1016/S1470-2045(17)30155-9.
 12. Olar A, Wani KM, Wilson CD, et al. Global epigenetic profiling identifies methylation

- subgroups associated with recurrence-free survival in meningioma. *Acta Neuropathol.* 2017;133(3):431-444. doi:10.1007/s00401-017-1678-x.
13. Reuss DE, Piro RM, Jones DTW, et al. Secretory meningiomas are defined by combined KLF4 K409Q and TRAF7 mutations. *Acta Neuropathol.* 2013;125(3):351-358. doi:10.1007/s00401-013-1093-x.
 14. Lee Y, Liu J, Patel S, et al. Genomic landscape of meningiomas. *Brain Pathol.* 2010;20(4):751-762. doi:10.1111/j.1750-3639.2009.00356.x.
 15. Claus EB, Park PJ, Carroll R, Chan J, Black PM. Specific genes expressed in association with progesterone receptors in meningioma. *Cancer Res.* 2008;68(1):314-322. doi:10.1158/0008-5472.CAN-07-1796.
 16. Hovestadt V, Zapatka M. conumee: Enhanced copy-number variation analysis using Illumina 450k methylation arrays. R package version 0.99. *R Packag.* 2015.
 17. Aizer AA, Abedalthagafi M, Linda Bi W, et al. A prognostic cytogenetic scoring system to guide the adjuvant management of patients with atypical meningioma. *Neuro Oncol.* 2016;18(2):269-274. doi:10.1093/neuonc/nov177.
 18. Heagerty PJ, Lumley T, Pepe MS. Time-dependent ROC curves for censored survival data and a diagnostic marker. *Biometrics.* 2000;56(2):337-344. doi:10.1111/j.0006-341X.2000.00337.x.
 19. Huang DW, Sherman BT, Tan Q, et al. The DAVID Gene Functional Classification Tool: A novel biological module-centric algorithm to functionally analyze large gene lists. *Genome Biol.* 2007;8(9):R183. doi:10.1186/gb-2007-8-9-r183.
 20. Chambless LE, Diao G. Estimation of time-dependent area under the ROC curve for long-term risk prediction. *Stat Med.* 2006;25(20):3474-3486. doi:10.1002/sim.2299.

21. Grambsch PM, Therneau TM. Proportional Hazards Tests and Diagnostics Based on Weighted Residuals Published by : Biometrika Trust Stable URL : <http://www.jstor.org/stable/2337123>. *Biometrika*. 2008;81(3):515-526.
22. Harmancl AS, Youngblood MW, Clark VE, et al. Integrated genomic analyses of de novo pathways underlying atypical meningiomas. *Nat Commun*. 2017;8:14433. doi:10.1038/ncomms14433.
23. Balachandran VP, Gonen M, Smith JJ, DeMatteo RP. Nomograms in oncology: more than meets the eye. *Lancet Oncol*. 2015;16(4):e173-80. doi:10.1016/S1470-2045(14)71116-7.
24. McShane LM, Cavenagh MM, Lively TG, et al. Criteria for the use of omics-based predictors in clinical trials. *Nature*. 2013;502(7471):317-320. doi:10.1038/nature12564.
25. Sproul D, Meehan RR. Genomic insights into cancer-associated aberrant CpG island hypermethylation. *Brief Funct Genomics*. 2013;12(3):174-190. doi:10.1093/bfgp/els063.
26. Huang RY, Bi WL, Weller M, et al. Proposed Response Assessment and Endpoints for Meningioma Clinical Trials: Report from the Response Assessment in Neuro-Oncology (RANO) Working Group. *Neuro Oncol*. August 2018. doi:10.1093/neuonc/noy137.

FIGURE LEGENDS

Figure 1. Comparison of grade-based and methylome-based recurrence-free survival predictor performance. Data presented are time-dependent receiver operating characteristic curve (ROC) and average area under the curve (AUC) as well as Δ AUC (DAUC) with 95%CI using 10,000 bootstrap resampling validation approach for methylome-based and grade-based predictors in the (A) first validation cohort and (B) second validation cohort (C) third validation cohort, (D) combined validation cohorts.

Figure 2. Recurrence-free survival (RFS) analysis of the first validation cohort (A), second validation cohort (B), and (C) third validation cohort using the 5y methylome-based RFS predictor, based on separation into distinct risk groups by median.

Figure 3. Frequency of copy number alterations across the genome stratified in risk groups according to the methylome-based predictor. Groups correspond to same groups seen in Figure 2.


Figure 4. Recurrence-free survival (RFS) analysis of all WHO grade I (A), WHO grade II (B) and WHO grade III (C) tumors in all three validation cohorts.

Figure 5. Nomogram to predict 5-year recurrence risk in meningiomas (A) Total points generated from scoring of methylome-based RFS predictor, WHO grade, and Simpson grade are tallied in the calculator and correlated to 5-year RFS in the nomogram in the Risk


Nomogram. (B) Time-dependent average AUC with 95%CI as well as Δ AUC (DAUC) with 95%CI using 10,000 bootstrap resampling validation approach generated for the meningioma recurrence score and a nomogram using clinical factors alone in the first and third validation cohorts as well as both combined validation cohorts. (C) Calibration curve of the nomogram to predict recurrence-free survival at 5 years in the combined validation cohort. The observed recurrence-free survival is plotted on the y-axis and nomogram predicted probability is plotted on the x-axis.

Figure 6. The meningioma recurrence score identifies two risk groups (high risk versus low risk) that may help individualize adjuvant management decisions such as the need for radiation therapy in patients with meningiomas. Patient tumor samples can be interrogated for DNA methylation profile of selected 9529 probes and a 5-year methylome-based RFS predictor score is generated. This score is combined with tumor WHO grade and Simpson grade in the meningioma recurrence score to develop an individualized probability of 5-year recurrence-free survival.


5-Year ROC




A. First Validation Cohort







B. Second Validation Cohort




C. Third Validation Cohort









A



B




C



Postop meningioma resection



- 1) Methylome-based predictor
- 2) Meningioma Recurrence Score



Low risk	201	192	183	179	175	169
High risk	200	172	131	100	82	69

High risk

Low risk

Consider escalation of
adjuvant therapy and
short-interval active
surveillance

Consider de-escalation
of adjuvant therapy and
regular active
surveillance